



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, A61K 31/70	A1	(11) International Publication Number: WO 99/45776 (43) International Publication Date: 16 September 1999 (16.09.99)
<p>(21) International Application Number: PCT/US98/24218</p> <p>(22) International Filing Date: 13 November 1998 (13.11.98)</p> <p>(30) Priority Data: 09/037,895 10 March 1998 (10.03.98) US 09/123,800 28 July 1998 (28.07.98) US</p> <p>(71) Applicant: MILKHAUS LABORATORY, INC. [US/US]; Larry Hill Road, Corner of Westfall, R.D. 1 Delanson, Delanson, NY 12053 (US).</p> <p>(72) Inventors: McMICHAEL, John; Corner of Westfall & Larry Hill Road, R.D.1 Delanson, Delanson, NY 12053 (US). ALLEN, Michael; 1149 38th Street, Sacramento, CA 95816 (US).</p> <p>(74) Agent: SHARP, Jeffrey, S.; Marshall, O'Toole, Gerstein, Murray & Borun, 6300 Sears Tower, 233 South Wacker Drive, Chicago, IL 60606-6402 (US).</p>		<p>(81) Designated States: AU, CA, IL, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: TREATMENT OF RESPIRATORY DISEASE AND OTITIS MEDIA</p> <p>(57) Abstract</p> <p>Methods for treating symptoms of respiratory disease and otitis media in a patient are presented. Methods comprise administering an effective amount of DNA to a subject in a manner so as not to effect gene transfer.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TREATMENT OF RESPIRATORY DISEASE AND OTITIS MEDIA

5

10

FIELD OF THE INVENTION

The present invention relates to methods for treatment of pulmonary disorders and otitis media.

BACKGROUND OF THE INVENTION

15

The present invention provides methods for treatment of pulmonary diseases. Such diseases, including cystic fibrosis, emphysema, chronic bronchitis, sinusitis, and the common cold, have in common bronchial or sinus congestion, production of large amounts of sputum, and the possibility of secondary bacterial infection requiring antibiotic therapy. The most serious

20 of those diseases is cystic fibrosis, a genetic disorder of exocrine function characterized by abnormally viscous mucus secretions leading to chronic pulmonary obstruction, pancreatic insufficiency and elevated sweat sodium and chloride levels. Cystic fibrosis is often fatal. The viscosity of sputum produced by cystic fibrosis patients is thought to be due to its high content of

25 DNA. Diseases such as bronchitis, emphysema, sinusitis, and the common cold are generally less severe than cystic fibrosis, but those diseases also may result in production of large amounts of sputum. Still other pulmonary diseases include mucositis (inflammation of the mucosal membranes) which is frequently

associated with radiation therapy and which is characterized by production of a thick water deficient mucous which is difficult for the subject to eliminate.

Other pulmonary diseases include chronic obstructive pulmonary diseases (COPDs) which share the common feature of chronic expiratory
5 airflow limitation i.e., persistent slowing of the rate at which exhalation can be achieved. Common COPDs include chronic bronchitis, emphysema and asbestosis and are characterized by respiratory distress but not associated with aberrant mucous accumulation. Cigarette smoke is the most common cause of COPDs which are also associated with exposure to respirable dusts particularly
10 in workplace environments of those engaged in occupations such as gold and coal mining, textile manufacturing and cement and steel making.

As with cystic fibrosis, other pulmonary diseases frequently lead to secondary bacterial infections. Treatment of pulmonary diseases generally requires antibiotic therapy which is frequently ineffective. Recently, however,
15 cystic fibrosis has been treated using DNase. The rationale for such therapy is that degrading DNA in sputum reduces the viscosity of the sputum and results in an increased ability of the patient to evacuate sputum from the lungs and nasal passages.

Acute otitis media is a bacterial or viral infection in the middle
20 ear which is usually secondary to upper respiratory tract infections and is most common in children. Microorganisms may migrate from the nasopharynx to the middle ear over the surface of the eustachian tube's mucous membrane or by propagating in the lamina propria of the mucous membrane as a spreading cellulitis or thrombophlebitis. Pain and hearing loss are the most common
25 presenting complaints although fever, nausea, vomiting and diarrhea may occur in young children. Therapy for acute otitis media includes analgesics, decongestants and antibiotics. In addition, topical vasoconstrictors may be administered into the nasal cavity to improve eustachian tube function. Further,

systemic sympathomimetic amines such as ephedrine sulfate may also be administered.

Serous otitis media (secretory otitis media) is an effusion in the middle ear resulting from incomplete resolution of acute otitis media or
5 obstruction of the eustachian tube. Traditional therapy includes a trial of antibiotic therapy in case of bacterial infection. Such antibiotic therapy is effective in relieving eustachian tube obstruction due to bacterial infection and in sterilizing the middle ear. Systemic sympathomimetic amines may also improve eustachian tube function by their vasoconstrictive effects and
10 antihistamines may relieve eustachian tube obstruction in allergic patients. Surgical therapies include myringotomy for aspiration of the fluid and for insertion of a tympanostomy tube which allows ventilation of the middle ear and ameliorates the eustachian tube obstruction. Alternatively, the middle ear may be temporarily ventilated with the Valsalva maneuver or politzeration.

15 Despite the efficacy of these approaches there remains a desire to avoid surgical intervention in cases of otitis media. Moreover, there exists a growing concern that the widespread use of antibiotics for treatment of otitis media in children promotes the development of antibiotic resistant bacteria. Accordingly, there remains a desire in the art for improved treatment of
20 conditions associated with upper respiratory infections and pulmonary disorders including otitis media.

SUMMARY OF THE INVENTION

The present invention provides methods for treating respiratory
25 illness. Specifically, the invention provides methods for treating symptoms of respiratory distress not associated with aberrant mucous accumulation in a patient, comprising the step of administering in a manner so as not to effect gene transfer an effective amount of DNA in a pharmaceutically-acceptable vehicle to a patient having a disease characterized by respiratory distress not

associated with aberrant mucous accumulation including but not limited to diseases such as chronic obstructive pulmonary disease including bronchitis, emphysema and asbestosis as well as asthma.

The invention further provides methods for relieving respiratory
5 congestion in a patient as a result of overproduction of viscous mucus or sputum lodged in the patients's respiratory tract due to conditions including mucositis such as caused by radiation comprising the steps of administering in a manner so as not to effect gene transfer a therapeutically effective amount of DNA in a pharmaceutically-acceptable vehicle to a patient having a disease characterized by
10 respiratory congestion, wherein said respiratory congestion is a result of an overproduction of viscous mucus or sputum lodged in said patients's respiratory tract, and wherein said method results in the reduced viscosity of said mucus or said sputum such that there is an increase of production and a reduced accumulation of mucus in said patient's respiratory tract.

15 Methods of the invention comprise administration to a patient suffering from respiratory distress an effective amount of DNA. The DNA is preferably provided in an amount ranging from about 0.00012 mg to about 0.003 mg and is preferably formulated in a liquid vehicle and provided at a concentration of approximately 0.0006 mg as single drops. A preferred route
20 of administration is sublingual, but other routes, such as subcutaneous, intravenous, intramuscular, and intrathecal are expected to work. DNA for use in the present invention may be prokaryotic DNA or eukaryotic DNA and may be formulated in a number of pharmaceutically-acceptable vehicles, including water, saline, albumin, and dextrose.

25 The present invention also provides methods for treating symptoms of otitis media in a patient, comprising the step of administering in a manner so as not to effect gene transfer an effective amount of a polynucleic acid which is preferably DNA in a pharmaceutically-acceptable vehicle to a patient having otitis media.

Methods of the invention comprise administration to a patient suffering from otitis media including acute otitis media, serous otitis media and chronic otitis media an effective amount of a polynucleic acid which is preferably DNA. The polynucleic acid may be selected from the group consisting of single-stranded and double-stranded DNA and RNA and includes natural polynucleic acids as well as synthetic nucleic acids such as poly-dT. The preferred polynucleic acid for use according to the invention is double-stranded DNA which is preferably provided in an amount ranging from about 0.00012 mg to about 0.003 mg and is preferably formulated in a liquid vehicle and provided at a concentration of approximately 0.0006 mg as single drops. A preferred route of administration is sublingual, but other routes, such as subcutaneous, intravenous, intramuscular, and intrathecal are expected to work. In addition, topical administration in the form of ear drops is also expected to be useful according to the invention. DNA for use in the present invention may be prokaryotic DNA or eukaryotic DNA including DNA from sources such as calf thymus, *E. coli* and salmon testicles. The DNA may be formulated in a number of pharmaceutically-acceptable vehicles, including water, saline, albumin, and dextrose.

Additional aspects and advantages of the invention will become apparent upon consideration of the following detailed description thereof.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides methods for treating patients with symptoms of respiratory distress not associated with aberrant mucous accumulation in a patient including symptoms resulting from chronic obstructive pulmonary diseases such as chronic bronchitis, emphysema, by administering to such patients a small amount of DNA in a manner so as not to effect gene transfer. Methods of the invention are also useful for treating pulmonary congestion in patients having any disease in which mucus production

is a symptom and are especially effective in treating diseases wherein viscous mucus or sputum is produced and becomes lodged in a patient's respiratory tract. In those cases, methods of the invention reduce production of DNA in a patient's mucus secretions and thereby render mucus less viscous, allowing
5 for increased production away from the respiratory tract.

Methods according to the invention for treating pulmonary congestion have been tested in clinical trials with human patients having various respiratory disorders, including cystic fibrosis, bronchitis, and emphysema using calf thymus DNA (Sigma, St. Louis). In each case, patients are
10 administered sublingual drops of DNA at a concentration of about 0.0006 mg DNA per drop. No other therapy was conducted in any patient during the course of DNA therapy. As noted below, all patients tested showed improvement in mucus production (*i.e.* sputum was easier to dislodge) from the respiratory tract. In addition, sputum was less viscous as compared to
15 pretreatment levels. Reduced sputum viscosity leads to increased patient comfort, increased ability of the patient to breathe, and reduced risk of secondary bacterial infection. No adverse side effects were observed in any patients. Drops of DNA may be administered at the appropriate concentration in doses of 1 to 10 drops per day as required by the patient.

20 The present invention also provides methods for treating patients with symptoms of otitis media by administering to such patients a small amount of DNA in a manner so as not to effect gene transfer. Methods of the invention are also useful for treating upper respiratory infections and pulmonary disorders including but not limited to those involving congestion in patients having any
25 disease in which mucus production is a symptom and are especially effective in treating diseases wherein viscous mucus or sputum is produced and becomes lodged in a patient's respiratory tract. In those cases, methods of the invention reduce production of DNA in a patient's mucus secretions and thereby render

mucus less viscous, allowing for increased production away from the respiratory tract.

Methods according to the invention for treating otitis media have been tested in clinical trials with human patients using salmon or calf thymus DNA. In each case, patients are administered sublingual drops of DNA at a concentration of about 0.0006 mg DNA per drop. In general, no other therapy was conducted in any patient during the course of DNA therapy. No adverse side effects were observed in any patients. Drops of DNA may be administered at the appropriate concentration in doses of 1 to 10 drops per day as required by the patient. For each Example below, salmon testicle or calf thymus DNA (Sigma) was used.

The following Examples illustrate the methods of the invention with respect to treatment of pulmonary diseases and in particular with respect to the preferred methods of treating otitis media. Numerous improvements and further aspects of the invention are apparent to the skilled artisan upon consideration of the Examples which follow.

EXAMPLE I

Twenty-three year-old twin brothers presented with cystic fibrosis. Each had a history of hospitalizations for lung clearance and secondary infections diagnosed as being associated with their cystic fibrosis. Each patient began therapy with 1-2 drops (0.0006 mg/drop) of DNA sublingually per day. For almost two years since beginning DNA therapy, neither patient has been hospitalized. In addition, follow-up evaluations by physicians revealed a 30-45% increase in airflow in each patient. Moreover, forced vital capacity, a common measure of lung capacity and the extent of mucus clearance in the lungs, increased from 60-90%. Finally, each of the brothers has gained weight and has shown increased expectoration.

After approximately one year of therapy, one of the brothers stopped taking the DNA drops. His condition steadily worsened as a result, with increased mucus viscosity, decreased forced vital capacity and reduced expectoration. That patient then began taking drops of DNA at the prescribed
5 dose and immediately improved to the condition he was in prior to the time at which he stopped taking the drops.

EXAMPLE II

A 64-year-old female patient who suffered from emphysema and
10 bronchitis, as diagnosed by her physician, was placed on a dose of 1 drop per day (0.0006 mg/drop) of DNA sublingually. Within one week, a follow-up evaluation revealed that her mucus production was less viscous and expectoration was increased.

EXAMPLE III

A 25-year-old female diagnosed with chronic upper respiratory
15 illness was treated with methods according to the invention. Previous antibiotic therapy was unsuccessful in treating her condition. She began with 1 drop of DNA (0.0006 mg/drop) sublingually four times per day. Within one day, she
20 experienced an increase in expectoration and, after three days she was able to discontinue treatment, having been completely relieved of congestion. She has remained symptom free.

EXAMPLE IV

A 32-year-old female nurse presented with a severe upper
25 respiratory infection and unproductive respiratory congestion. She was placed on 1 drop of DNA (0.0006 mg/drop) four times per day. Her congestion began to break up almost immediately. Expectoration was improved and the patient's illness resolved after 4.5 days and no congestion recurred.

EXAMPLE V

A 63-year-old woman presented with chronic sinusitis. Four drops of DNA per day were administered. After 3 months, the patient's mucus had thinned and her cough was more productive.

5

EXAMPLE VI

A 37-year-old female presented with unresolved respiratory congestion. Traditional therapy, including expectorants, failed to improve her condition. The patient was then prescribed four drops of DNA (0.0006 mg/drop) per day. After one day of treatment, her congestion was more productive and sinus drainage had begun where none was present prior to treatment according to the invention.

10

EXAMPLE VII

A 40-year-old woman with unproductive upper respiratory congestion was placed on 4 drops of DNA (0.0006 mg/drop) per day. Her congestion was more productive after one day and she continued to expectorate freely. In this case, therapy was supplemented with an over-the-counter expectorant.

15

20

EXAMPLE VIII

A 38-year-old woman with acute and chronic respiratory disease due to exposure to toxic corrosive materials was treated with methods according to the invention. Prior to such treatment, symptoms, including chronic rhinorrhea, chest congestion and chronic respiratory infections were treated with numerous courses of antibiotics without success. The patient began treatment with 0.5 cc Q.I.D. daily and was instructed to administer treatment up to 5-6 times daily if necessary.

25

Upon commencing treatment according to the invention, the patient was able to produce sputum almost immediately. Continued treatment has alleviated symptoms of chronic respiratory illness.

5

EXAMPLE IX

A 58-year-old woman diagnosed with a childhood history of asthma and persistent adult rhinitis and sinusitis presented for treatment. Physical examination indicated clear rhinorrhea, and 3+ red throat. Nasal spray and prednisone were prescribed for 7 days. That course of treatment
10 resulted in mild improvement. However, the patient's cough was still unproductive. Therapy according to the invention was begun at 0.5 cc Q.I.D. Within 48 hours, the patient showed improvement in the form of a productive cough and sinus drainage.

15

EXAMPLE X

A 48-year-old woman with chronic sinusitis and bronchitis characterized by chronic head congestion, nasal obstruction, and coughing presented for treatment according to the invention. The patient was treated according to the invention with one drop per day of DNA (0.0006 mg/drop).
20 Treatment resulted in an overt increase in sinus and chest drainage. Upon cessation of treatment according to the invention, the patient's condition regressed. Beginning therapy again caused a similar increase in drainage and relief of congestion as seen previously with treatment according to the invention.

25

The following examples report the results of treatment of subjects suffering from radiation induced mucositis with the DNA containing compositions of the invention.

EXAMPLE XI

According to this example, a subject suffering from radiation induced mucositis was treated with one drop of DNA (0.0006 mg/drop) sublingually four times per day. The subject experienced a 50% improvement with phlegm thickness and had less cough. Experimentation by the subject with dosage frequency revealed that administration of one drop alone was insufficient but that administration of three to four drops per day appeared to be optimal.

10

EXAMPLE XII

According to this example, a subject suffering from radiation induced mucositis was treated with one drop of DNA (0.0006 mg/drop) sublingually four times per day. While treatment with four drops per day did not provide subjective improvement an increase in dosage to ten drops per day may have resulted in less phlegm. The subject discontinued administration of DNA but restarted use later and reported thinning of phlegm. The formulation was later modified to include 2 units of streptolysin O per drop although it could not be determined if incorporation of streptolysin O improved the therapeutic results.

15
20**EXAMPLE XIII**

According to this example, a subject suffering from radiation induced mucositis was treated with one drop of DNA (0.0006 mg/drop) sublingually four times per day with the result of a 50% improvement in phlegm thickness. In addition the subject noted that her sense of taste improved from nonexistent to normal.

25

The following examples report the results of treatment of three patients suffering with mild to moderate chronic obstructive pulmonary disease not characterized by aberrant mucous accumulation who were successfully

treated with DNA containing compositions according to the methods of the invention.

EXAMPLE XIV

5 A 67 year-old male former smoker with a medical history of
gout, hypertension, peptic ulcer and chronic obstructive pulmonary disease
presented with shortness of breath during high humidity, walking up a half
flight of stairs, walking in the yard and at night laying flat in bed. The subject
suffered from minimal phlegm production which was white in color. The
10 subject was being treated with allopurinol, Pepcid (famotidine), Slobid
(theophylline), Calan (verapamil HCl), Accupril (quanapril HCl) and Albuterol
Inhaler. A pre-study office spirometry showed moderate COPD with an Fev1
% of 51.

 The subject was treated with 1 drop of DNA (0.0006 mg/drop)
15 sublingually four times per day. After fourteen days of treatment the subject
reported that his overall dyspnea had improved from a subjective rating of a 10
to a 4. He was able to walk at the mall without shortness of breath where
previously, he had to stop. A spirometry on day 16 showed no change but
three months later with continued treatment according to the invention could
20 ascend 13 steps where prior to treatment he had been unable to ascend only half
as many steps without dyspnea. The subject was also able to decrease Albuterol
administration from daily to 2-3 times weekly and eventually to once in four
weeks and discontinue use of Slobid. The subjects wife reported that the
subject's sleep is more restful and that she no longer hears wheezing at night.

25

EXAMPLE XV

 A 71 year-old female with a medical history of hypertension,
myocardial infarction, renal insufficiency, hiatal hernia, spinal stenosis,
hyperlipidemia and chronic obstructive pulmonary disease presented with

shortness of breath while cooking meals, walking 17 steps, carrying laundry, vacuuming, making her bed, walking to the car, and in the mall. She also complained of minimal phlegm. She was undergoing treatment with medications including Cardizem CD (diltiazem HCl), Vasotec (enalaprilat),
5 Zocor (simvastatin), Ogen (estropipate), Zantac (ranitidine HCl), Toprol (metoprolol succinate), Nitroglycerine patch, LorTab (hydrocodone bitartrate and aspirin), and a sleep agent as required. Upon examination, she had mild anterior wheezing and a pre-study office spirometry showed an Fev1 of 70.

The subject was treated with 1 drop of DNA (0.0006 mg/drop)
10 sublingually four times per day. After seven days of treatment the subject reported no improvement but fourteen days of treatment reported that she could walk in the mall without shortness of breath and was vacuuming and making her bed without needing to stop and rest. A repeat spirometry after fourteen days showed an Fev1 % of 78, an 11% improvement from the pre-study result.
15 The subject's condition continued to improve except when she decreased the treatment schedule to once per day and her shortness of breath returned. After increasing back to treatment four times daily her dyspnea resolved to the extent that she was able to discontinue her use of a Serevent (Glaxo) aerosol inhaler after four months.

20

EXAMPLE XVI

A 76 year-old female with a medical history of hypertension, arrhythmia, hypercholesterolemia, chronic obstructive pulmonary disease (for at least ten years) and anxiety presented with dyspnea after climbing one flight
25 of stairs, exertional dyspnea and cough and with minimal phlegm. The subject was being treated with Normodyne (labetalol HCl), Procardia (nifedipine), Persantine (dipyridamole), Zocor (simvastatin), Atrovent Inhaler (ipratropium bromide) and Xanax (alprazolam). Upon examination, she had moderately decreased lung sounds with normal blood pressure. A spirometry conducted ten

years previously showed an Fev1 % of 73 (normal) with diminished mid flow rates suggesting early COPD.

The subject was treated with 1 drop of DNA (0.0006 mg/drop) sublingually twice daily and after one month of treatment had less coughing and diminished wheezing at home when in bed. A spirometry after almost two months of treatment showed an Fev1 % of 65. The subject continued to report subjective improvement and stopped administration of Atrovent. After four months wheezing was nearly gone and her cough was less than prior to treatment according to the invention.

10

EXAMPLE XVII

According to this example, several asthma patients were treated by daily administration of at least one drop of DNA (0.0006 mg/drop) derived from either salmon sperm or bovine sources. Follow-up evaluation of those subjects showed decreased viscosity and volume of sputum. In addition, the salmon sperm DNA was found to have therapeutic activity equivalent to that of the bovine derived DNA.

15

EXAMPLE XVIII

According to this example, a 56 year old non-smoker with chronic obstructive pulmonary disease/emphysema secondary to asbestosis and total disability due to pulmonary insufficiency was treated by sublingual administration of at least one drop of DNA (0.0006 mg/drop) four times daily. After a few weeks of treatment the subject reported feeling "dramatically better" and "not out of breath." The subject has since reduced the frequency of treatment to one drop daily.

20

25

EXAMPLE XIX

According to this example, the sublingual administration of one drop of DNA (0.0006 mg/drop) provided almost immediate relief of symptoms of respiratory disorders caused by chemical or environmental sensitivities. The therapy was tested on at least a dozen individuals including children and adults. and was successful in all cases.

In the following examples, administration of DNA derived from either salmon testicles or calf thymus (Sigma) was found to be useful in treatment of otitis media with rapid and reproducible responses.

EXAMPLE XX

According to this example, a five year old female presented with severe recurrent otitis media in the right ear with bulging of the tympanic membrane. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) four times daily for seven days. When the subject was rechecked two days later the mother reported the child's temperament and energy improved the first evening. She went to school the next day. On exam, she had an injected tympanic membrane, but the bulging was gone. Significantly, this subject has been treated for OM numerous times in the past with antibiotics.

Roughly nine days after termination of treatment according to the invention the subject developed recurrent pain in the left ear with a fever of 101° F. Her mother did not contact the physician but administered the DNA composition of the invention again with success. She went to school the next day. The patient presented again April 2, 1998 with recurrent otitis media in the right ear. She was again treated by sublingual administration of the DNA composition of the invention resulting in rapid sense of pain relief, temperature resolution and improved overall well being.

EXAMPLE XXI

According to this example, a nine year old female presented with a plugged feeling in the right ear, fever of 100° F and minimal pain. A right
5 otitis media was diagnosed. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) four times daily. The first night of treatment, the child slept well, the pain left and she went to school the next day. Four days later the redness and fluid were less. The mother reported that usually with otitis media and antibiotic treatment, her child had
10 2-3 restless nights and usually missed 2-3 days of school. In the case of treatment according to the invention she went to school the next day.

EXAMPLE XXII

According to this example, a five year old female presented with
15 acute otitis media in which she had a fever, was whining and was restless. The mother gave her one dose of a plant-derived homeopathic remedy in the late afternoon. It helped some, but later that night, severe pain recurred. The subject was treated with a single sublingual administration of one drop of DNA (0.0006 mg/drop). No further doses were given. The child went to pre-school
20 the next day. The ear was checked several times the next three weeks and gradually it returned to normal.

EXAMPLE XXIII

According to this example, a seventeen month old male presented
25 with recurrent serous otitis media. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) four times daily for one month. There occurred complete resolution of the fluid.

EXAMPLE XXIV

According to this example, a three year old presented with recurrent serous otitis media. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) hourly until pain was eliminated. The subject was then treated four times daily for one week and had a complete resolution of symptoms.

EXAMPLE XXV

According to this example, a two year old female presented with bilateral otitis media with pain. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) hourly until the pain was relieved and then one drop four times daily for one week and had a complete resolution of symptoms.

EXAMPLE XXVI

According to this example, a ten month old male presented with bilateral serous otitis media and eustachian tube dysfunction. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) four times daily for one month. Reevaluation after one month showed normal eustachian tubes.

EXAMPLE XXVII

According to this example, a fourteen month old male presented with recurrent bilateral serous otitis media for which surgery to insert tympanostomy tubes into the eustachian tubes was originally recommended. The subject was treated with sublingual administration of one drop of DNA (0.0006 mg/drop) four times daily. Evaluation one and two months later revealed incomplete resolution of fluid but evaluation three months later revealed complete elimination of fluid. Five months after initiation of therapy

bilateral otitis media recurred but resumption of therapy with the DNA drops resulted in a complete resolution.

The invention has been described in terms of its preferred
5 embodiments and is only intended to be limited by the scope of the following claims.

WHAT IS CLAIMED IS:

1. A method for treating symptoms of respiratory distress not associated with aberrant mucous accumulation in a patient, comprising the step of:
5 administering in a manner so as not to effect gene transfer an effective amount of DNA in a pharmaceutically-acceptable vehicle to a patient having a disease characterized by respiratory distress not associated with aberrant mucous accumulation.
- 10 2. The method according to claim 1, wherein said DNA is administered sublingually in the form of a liquid drop.
3. The method according to claim 1, wherein said disease is selected from the group consisting of chronic bronchitis, emphysema and asthma.
15
4. The method according to claim 1, wherein said disease is asthma.
5. The method according to claim 1, wherein said vehicle is
20 selected from the group consisting of water, saline, albumin, or dextrose.
6. The method according to claim 1, wherein said effective amount of DNA is from about 0.00012 mg to about 0.003 mg DNA.
- 25 7. The method according to claim 1, wherein said effective amount of DNA is about 0.0006 mg of DNA.
8. The method according to claim 1 wherein said patient is a human.
30

9. The method of claim 1 wherein the DNA is administered by a route selected from the group consisting of sublingual, subcutaneous, intravenous, intramuscular and intrathecal administration.

5 10. A method for relieving respiratory congestion associated with mucositis in a patient, comprising the steps of:

 administering in a manner so as not to effect gene transfer a therapeutically effective amount of DNA in a pharmaceutically-acceptable vehicle to a patient having a disease characterized by respiratory congestion, wherein said
10 respiratory congestion is a result of an overproduction of viscous mucus or sputum lodged in said patients's respiratory tract, and wherein said method results in the reduced viscosity of said mucus or said sputum such that there is an increase of production and a reduced accumulation of mucus in said patient's respiratory tract.

15

 11. The method according to claim 10, wherein said DNA is administered sublingually in the form of a liquid drop.

 12. The method according to claim 10, wherein said vehicle is
20 selected from the group consisting of water, saline, albumin, or dextrose.

 13. The method according to claim 10, wherein said effective amount of DNA is from about 0.00012 mg to about 0.003 mg DNA.

25 14. The method according to claim 10, wherein said effective amount of DNA is about 0.0006 mg of DNA.

 15. The method according to claim 10 wherein said patient is a human.

30

16. The method of claim 10 wherein the DNA is administered by a route selected from the group consisting of sublingual, subcutaneous, intravenous, intramuscular and intrathecal administration

5 17. A method for treating symptoms of otitis media, comprising the step of:
administering in a manner so as not to effect gene transfer an effective amount of DNA in a pharmaceutically-acceptable vehicle to a patient having otitis media.

10

18. The method according to claim 17 wherein said DNA is administered sublingually in the form of a liquid drop.

19. The method according to claim 17 wherein said vehicle is
15 selected from the group consisting of water, saline, albumin, or dextrose.

20. The method according to claim 17 wherein said effective amount of DNA is from about 0.00012 mg to about 0.003 mg DNA.

20 21. The method according to claim 17 wherein said effective amount of DNA is about 0.0006 mg of DNA.

22. The method according to claim 17 wherein said patient is
a human.

25

23. The method of claim 17 wherein the DNA is administered by a route selected from the group consisting of sublingual, subcutaneous, intravenous, intramuscular, intrathecal and topical administration.

24. The method of claim 17 wherein the DNA is administered topically in the form of eardrops.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/24218

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A01N 43/04; A61K 31/70

US CL :514/44, 2, 47, 50; 536/22.1; 424/9.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44, 2, 47, 50; 536/22.1; 424/9.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, medline, caplus, biosis, wpids

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P -- Y,P	US 5,726,160 A (MCMICHAEL, J.) 10 March 1998, see claims 1-7.	1-9 ----- 10-24
X -- Y	WO 9632138 A1 (MILKHAUS LABORTORY) 17 October 1996, see claims.	1-9 ----- 10-24



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

24 DECEMBER 1998

Date of mailing of the international search report

15 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Michael C. Wilson

Telephone No. (703) 308-0120